

# A partition of connected graphs

Gus Wiseman

Department of Mathematics, University of California, One Shields Ave., Davis, CA 95616

[gus@nafindex.com](mailto:gus@nafindex.com)

Published in The Electronic Journal of Combinatorics, Jan 7, 2005

Mathematics Subject Classifications: 05C30, 05C05

## Abstract

We define an algorithm  $k$  which takes a connected graph  $G$  on a totally ordered vertex set and returns an increasing tree  $R$  (which is not necessarily a subtree of  $G$ ). We characterize the set of graphs  $G$  such that  $k(G) = R$ . Because this set has a simple structure (it is isomorphic to a product of non-empty power sets), it is easy to evaluate certain graph invariants in terms of increasing trees. In particular, we prove that, up to sign, the coefficient of  $x^q$  in the chromatic polynomial  $\chi_G(x)$  is the number of increasing forests with  $q$  components that satisfy a condition that we call  $G$ -connectedness. We also find a bijection between increasing  $G$ -connected trees and broken circuit free subtrees of  $G$ .

We will work with finite labeled simple graphs. Usually we will identify a graph  $G$  with its edge set; this should not cause any serious ambiguities. If the vertex set is  $V$  then we say that  $G$  is a graph on  $V$ . A (spanning) subgraph  $Q$  of  $G$  is a graph with the same vertex set as  $G$  and a subset of the edges of  $G$ . The notation  $Q \subseteq G$  means  $Q$  is a subgraph of  $G$ . A rooted graph is a graph with a distinguished vertex called the root.

Define  $\text{link}(v, S)$  to be the set of all possible edges joining  $v$  to an element of  $S$  (so if  $v \notin S$ ,  $\text{link}(v, S)$  has  $|S|$  elements). If  $G$  is a graph on  $V$  and  $S \subseteq V$ , we define the restriction of  $G$  to  $S$ ,  $G|_S$ , to be the graph on  $S$  whose edge set consists of all edges of  $G$  with both ends in  $S$ .

We will use the symbols  $\pi$  and  $\sigma$  to denote set partitions. The notation  $\pi \vdash S$  means  $\pi$  is a set partition of the set  $S$ . The length (number of blocks) of  $\pi$  is denoted by  $\ell(\pi)$ . A set partition  $\sigma$  is called a refinement of a set partition  $\pi$  if every block of  $\sigma$  is contained in some block of  $\pi$ .

To each graph  $G$  on  $V$  there corresponds a set partition  $s(G)$  such that two vertices  $v, w \in V$  are in the same block of  $s(G)$  if and only if there is a path in  $G$  from  $v$  to  $w$ . Equivalently,  $s(G)$  is the maximal set partition of  $V$  whose blocks are connected. The restriction of  $G$  to a block of  $s(G)$  is called a component of  $G$ .

If  $G$  is a rooted connected graph on  $V$  with root  $r$ , we will call the set partition  $\pi = s(G|_{V-\{r\}})$  of  $V - \{r\}$  the depth-first partition of  $G$ . To obtain a connected subgraph

of a rooted connected graph  $G$  on  $V$ , we can choose, for each block  $\pi_i$  of  $\pi$ , a connected subgraph of  $G|_{\pi_i}$  and a nonempty set of edges (in  $G$ ) connecting  $r$  to  $\pi_i$ . In fact, every connected subgraph of  $G$  can be obtained in this way. Our Theorem 1 may be regarded as an iteration of this correspondence. The depth-first partition and this correspondence have been studied by Gessel [3].

A forest is a graph with no circuits. A tree is a connected forest. A basic property of trees is that there is a unique path (a sequence of distinct, adjacent vertices) between any two vertices. The distance between two vertices is defined to be the length of this path. In a rooted tree, the height of a vertex is defined to be its distance from the root. A vertex  $w$  is called a descendant of a vertex  $v$  (or  $v$  is called an ancestor of  $w$ ) if the heights of the vertices on the unique path from  $v$  to  $w$  are increasing (so in particular  $v$  is always a descendant of itself). We define the join of  $v$  and  $w$  to be their unique common ancestor on the unique path between them.

Let  $R$  be a rooted tree on the vertex set  $V$ , and let  $v \in V$ . We define  $\text{des}(v, R) \subseteq V$  to be the set of descendants of  $v$  (including  $v$ ). If  $v$  is not the root of  $R$ , we define  $\text{parent}(v, R) \in V$  to be the closest vertex to  $v$  in  $R$  which is not a descendant of  $v$ . A rooted tree is increasing (according to a total order on  $V$ ) if for each  $v \in V$  and  $w \in \text{des}(v, R)$  we have  $v \leq w$ . Consequently, the root of an increasing tree must be the smallest element of  $V$ .


**Definition 1** *Let  $R$  be a rooted tree on the totally ordered vertex set  $V$  with root  $r$ , and let  $v \in V - \{r\}$ . Define  $J(v, R) = \text{link}(\text{parent}(v, R), \text{des}(v, R))$ . If  $G$  is a graph on  $V$  and if for each  $v \in V - \{r\}$  we have  $J(v, R) \cap G \neq \emptyset$  then we say that  $R$  is  $G$ -connected.*

Note that the sets  $J(v, R)$  (as  $v$  ranges over  $V - \{r\}$ ) are disjoint. Also note that a  $G$ -connected tree need not be a subgraph of  $G$  and that  $G$  must be connected for any rooted tree to be  $G$ -connected.

**Definition 2** *For each connected graph  $G$  on a totally ordered vertex set  $V$ , define an increasing  $G$ -connected tree  $k(G)$  by the following algorithm:*

1. *Let  $H$  be an empty graph on  $V$ , and set  $S = V$ .*
2. *Let  $\pi$  be the depth-first partition of  $G|_S$  rooted at  $r$  = the smallest vertex in  $S$ . Add edges to  $H$  connecting  $r$  to the smallest vertex in each block of  $\pi$ .*
3. *For each block  $\pi_i$  of  $\pi$  with more than one element, return to step 2 with  $S = \pi_i$ .*
4. *Return  $k(G) = H$ .*

**Example 1** *The 6 increasing trees on  $V = \{1, 2, 3, 4\}$  are listed vertically. To the right of each increasing tree  $R$  are listed the subtrees  $T$  of the complete graph on  $V$  such that  $k(T) = R$  (we have omitted the 22 connected subgraphs which are not trees). The breaks are indicated by dotted lines (see Theorem 3).*



There is a different algorithm, called depth-first search, which produces subforests of  $G$ . Some enumerative applications of this algorithm have been studied by Gessel and Sagan [4]. A distinguishing difference between depth-first search and our algorithm is that depth-first search only follows the edges of  $G$ , whereas here we add edges connecting to the smallest vertex in each block of  $\pi$  regardless of whether these are edges of  $G$ . The algorithms are related in that if  $G$  is a connected graph and  $R$  is a depth-first search subtree of  $G$  then parts 2 and 3 of the next theorem hold (although the converse is not true).

**Theorem 1** *Let  $G$  be a connected graph on a totally ordered vertex set  $V$ , and let  $R$  be an increasing  $G$ -connected tree on  $V$ . Then the following are equivalent:*

1.  $k(G) = R$
2. *For each vertex  $v \in V$ ,  $G|_{\text{des}(v,R)}$  rooted at  $v$  is connected and has the same depth-first partition as  $R|_{\text{des}(v,R)}$  rooted at  $v$ .*
3. *For each non-root vertex  $v \in V - \{r\}$  there is a nonempty set  $E(v) \subseteq J(v, R)$  such that  $G = \bigcup_{v \in V - \{r\}} E(v)$ .*

*Proof.*  $1 \Leftrightarrow 2$  This follows easily from Definition 2.

$2 \Rightarrow 3$  Let  $E(v) = J(v, R) \cap G$ . We need to show that every edge of  $G$  lies in some  $E(v)$ . Let  $e \in G$  and let  $v < w$  be the vertices of  $e$ . We will show that  $w$  is a descendant of  $v$ . Suppose this is false, and let  $u$  be their join. Then  $e \in G|_{\text{des}(u, R)}$ , so  $v$  and  $w$  are in the same block of the depth first partition of  $G|_{\text{des}(u, R)}$ . This is a contradiction because they are in different blocks of the depth first partition of  $R|_{\text{des}(u, R)}$ . Now, since  $w$  is a descendant of  $v$ , there is a unique vertex  $z \in V$  (possibly equal to  $w$ ) such that  $\text{parent}(z) = v$  and  $w \in \text{des}(z)$ . Hence  $e \in J(z, R) \cap G$ .

$3 \Rightarrow 2$  This is certainly true if  $v$  (in part 2) is a leaf of  $R$  (its only descendant is itself). Let  $v \in V$  and suppose it is true for all  $w \in \text{des}(v, R) - \{v\}$ . Let  $\pi$  be the depth-first partition of  $R|_{\text{des}(v, R)}$ . Then  $G|_{\pi_i}$  is connected by the inductive hypothesis. Furthermore,  $G$  contains an edge connecting  $v$  to  $\pi_i$  because  $\pi_i$  contains a vertex  $w$  whose parent in  $R$  is  $v$  and  $J_G(w, R)$  consists of edges connecting  $v$  to  $\pi_i$ . Hence  $G|_{\text{des}(v, R)}$  is connected. Clearly  $\pi$  is a refinement of the depth-first partition of  $G|_{\text{des}(v, R)}$  (because  $G|_{\pi_i}$  is connected), so to show that they are equal we have only to show that if  $x$  and  $y$  are in different blocks of  $\pi$  then they are in different blocks of the depth-first partition of  $G$ . Let  $x < y \in V$  be in different blocks of  $\pi$ , and suppose  $G$  has an edge between  $x$  and  $y$ . Then  $y$  is a descendant of  $x$  in  $R$  because every edge of  $J_G(w, R)$  (for any  $w \in V$ ) connects a vertex to one of its descendants. This contradicts the fact that they are in different blocks of the depth-first partition of  $R|_{\text{des}(v, R)}$ .  $\square$

**Remark 1** *Actually the condition in Theorem 1 that  $R$  be  $G$ -connected is not necessary because if  $R$  is not  $G$ -connected then parts 1, 2 and 3 will be false.*

Some algebraic invariants of graphs can be simply expressed in terms of connected subgraphs. We can use the algorithm  $k$  to express such invariants in terms increasing trees. Moreover, Theorem 1 shows that the set  $k^{-1}(R)$  has a simple structure, as illustrated by the next theorem.

**Definition 3** *Let  $G$  be a connected graph on  $V$ . Define*

$$\eta^G(t) = \sum_{\substack{Q \subseteq G \\ \text{connected}}} t^{|Q|}$$

where  $|Q|$  denotes the number of edges in  $Q$ .

**Theorem 2**

$$\eta^G(t) = \sum_{\substack{R \\ \text{increasing} \\ G\text{-connected}}} \prod_{v \in V - \{r\}} [(1+t)^{|J(v, R) \cap G|} - 1]$$

*Proof.* We have

$$\eta^G(t) = \sum_{\substack{R \\ \text{increasing} \\ G\text{-connected}}} \sum_{\substack{Q \subseteq G \\ k(Q)=R}} t^{|Q|}$$

Now, the generating function for the cardinality of nonempty subsets of a set  $S$  is

$$f_S(x) = \sum_{\emptyset \neq T \subseteq S} x^{|T|} = (1+x)^{|S|} - 1$$

Hence from Theorem 1 part 3,

$$\sum_{\substack{Q \subseteq G \\ k(Q)=R}} t^{|Q|} = \sum_{\substack{Q = \bigcup_{v \in V - \{r\}} E(v) \\ \emptyset \neq E(v) \subseteq J(v, R) \cap G}} t^{|Q|} = \prod_{v \in V - \{r\}} f_{J(v, R) \cap G}(t)$$

from which the result follows.  $\square$

The chromatic polynomial  $\chi_G(x)$  of a graph  $G$  is a polynomial which evaluates to the number of proper colorings of  $G$  with  $x$  colors. The subgraph expansion of  $\chi_G(x)$  is

$$\chi_G(x) = \sum_{Q \subseteq G} (-1)^{|Q|} x^{c(Q)}$$

where  $c(Q)$  is the number of components of  $Q$ . See [1] for background on the chromatic polynomial.

We define an increasing  $G$ -connected forest  $R$  to be a forest where each component  $R|_{s(R)_i}$  is an increasing  $G|_{s(R)_i}$ -connected tree. For a graph  $G$ , let  $t(G)$  be the (integer) partition whose parts are the sizes of the blocks of  $s(G)$ . For background on the chromatic symmetric function  $X_G = X_G(x_1, x_2, \dots)$  of a graph  $G$ , see [5] and [6]. For background on the chromatic symmetric function in non-commuting variables  $Y_G = Y_G(x_1, x_2, \dots)$ , see [2].

**Corollary 1** *Let  $G$  be a graph on a totally ordered vertex set  $V$  with  $|V| = n$ .*

1. *The coefficient of  $(-1)^{n-1}x$  in the chromatic polynomial  $\chi_G(x)$  is the number of increasing  $G$ -connected trees.*
2. *The coefficient of  $(-1)^{n-q}x^q$  in the chromatic polynomial  $\chi_G(x)$  is the number of increasing  $G$ -connected forests with  $q$  components (or, equivalently, with  $n-q$  edges).*
3. *The coefficient of  $(-1)^{n-\ell(\lambda)}p_\lambda$  in the chromatic symmetric function  $X_G$  is the number of increasing  $G$ -connected forests  $R$  such that  $t(R) = \lambda$ .*
4. *The coefficient of  $(-1)^{n-\ell(\pi)}p_\pi$  in the chromatic symmetric function in non-commuting variables  $Y_G$  is the number of increasing  $G$ -connected forests  $R$  such that  $s(R) = \pi$ .*

*Proof.* 1. Let  $a^G$  be the coefficient of  $x$  in  $\chi_G(x)$ . From the subgraph expansion we have

$$a^G = \sum_{\substack{Q \subseteq G \\ \text{connected}}} (-1)^{|Q|} = \eta^G(-1) = \sum_{\substack{R \\ \text{increasing} \\ G\text{-connected}}} \prod_{v \in V - \{r\}} (-1)$$

We don't need to worry about  $0^0$  because the  $G$ -connectedness of  $R$  implies that  $J(v, R) \cap G$  is never empty.

4. We will prove part 4, the others being simple specializations. Let  $H_\pi^G$  be the number of increasing  $G$ -connected forests  $R$  such that  $s(R) = \pi$ , and let  $H^G$  be the number of increasing  $G$ -connected trees. Then using part 1 we have

$$H_\pi^G = \prod_{i=1}^{\ell(\pi)} H^{G|_{\pi_i}} = (-1)^{n-\ell(\pi)} \prod_{i=1}^{\ell(\pi)} \sum_{\substack{Q \subseteq G|_{\pi_i} \\ \text{connected}}} (-1)^{|Q|} \quad (1)$$

The subgraph expansion of  $Y_G$  is

$$Y_G = \sum_{Q \subseteq G} (-1)^{|Q|} p_{s(Q)}$$

Hence

$$Y_G = \sum_{\pi \vdash V} p_\pi \sum_{\substack{Q \subseteq G \\ s(Q) = \pi}} (-1)^{|Q|} = \sum_{\pi \vdash V} p_\pi \prod_{i=1}^{\ell(\pi)} \sum_{\substack{Q \subseteq G|_{\pi_i} \\ \text{connected}}} (-1)^{|Q|}$$

Substituting (1), we obtain the desired result.  $\square$

If  $G$  is a graph on a totally ordered vertex set  $V$ , we extend the ordering of the vertices to an ordering of the edges lexicographically. A broken circuit of  $H \subseteq G$  is a set of edges  $B \subseteq H$  such that there is some edge  $e \in G$ , smaller than every edge of  $B$ , such that  $B \cup e$  is a circuit. Note that  $B$  being a broken circuit of  $H$  depends both on  $H$  and  $G$ . If  $H \subseteq G$  contains no broken circuits then it is called broken circuit free. Note that if  $H$  contains a circuit then it also contains a broken circuit. Consequently, a broken circuit free subgraph is always a forest. If  $T \subseteq G$  is a subtree of  $G$  and the edge  $e \in G$ ,  $e \notin T$  is the smallest edge in the unique circuit in  $T \cup \{e\}$  then we will call  $e$  a break in  $T$ . Hence the set of breaks in a subtree  $T$  is in bijection with the set of broken circuits of  $T$ .

Whitney's Broken Circuit Theorem [7] shows that if  $G$  is a connected graph with  $n$  vertices, the coefficient of  $(-1)^{n-1}x$  in  $\chi_G(x)$  is the number of broken circuit free subtrees of  $G$ . Hence there should be a bijection between broken circuit free subtrees and increasing  $G$ -connected trees.

**Theorem 3** *Let  $V$  be a totally ordered vertex set with smallest element  $r$ , and let  $G$  be a connected graph on  $V$ . Let  $T \subseteq G$  be a subtree of  $G$ , and let  $R = k(T)$ . Let  $E(v)$  for  $v \in V - \{r\}$  be as in Theorem 1 part 3. Then  $E(v)$  contains only one element  $e(v)$  (otherwise  $T$  would have more than  $|V|-1$  edges so it could not be a tree). For  $v \in V - \{r\}$ , let  $d(v)$  be the set of elements of  $J(v, R) \cap G$  which are smaller than  $e(v)$ . Then the set of breaks in  $T$  is*

$$\bigcup_{v \in V - \{r\}} d(v)$$

*Proof.* Let  $J = \bigcup_{v \in V - \{r\}} J(v, R) \cap G$ . Since  $k(G)$  may be different from  $R$ ,  $J$  may be different from  $G$ . We will first show that if  $e \in G$  but  $e \notin J$  then  $e$  is not a break. Let  $v < w \in V$  be the vertices of  $e$ . Then  $w$  is not a descendant of  $v$  because otherwise we would have  $e \in J$ . Let  $u \in V$  be the join of  $v$  and  $w$  in  $R$ . Then Theorem 1 part 2 implies that  $u$  is also the join of  $v$  and  $w$  in  $T|_{\text{des}(u, R)}$  (rooted at  $u$ ). Therefore, the cycle created by adding  $e$  to  $T$  contains an edge connected to  $u$ . Since  $u < v < w$ ,  $e$  cannot be a break.

Now suppose  $e \in J(v, R) \cap G$  is smaller than  $e(v)$ . We will show that  $e$  is a break. Let  $H = T|_{\text{des}(v, R) \cup \text{parent}(v, R)}$ . Then  $\text{parent}(v, R)$  is the smallest vertex in the vertex set of  $H$ . Therefore,  $e$  is smaller than any other edge in  $H$ . Since  $H$  is a tree, adding  $e$  would create a unique circuit in  $H$ . Hence  $e$  is a break.

Now suppose  $e \in J(v, R) \cap G$  is larger than  $e(v)$ . Then, letting  $H$  be as before, we see that  $e(v)$  must belong to the circuit which  $e$  creates. But  $e(v)$  is smaller than  $e$ , so  $e$  cannot be a break.  $\square$

**Corollary 2** *The function*

$$f(R) = \bigcup_{v \in V - \{r\}} \min(J(v, R) \cap G)$$

*is a bijection between increasing  $G$ -connected trees and broken circuit free subtrees, and  $f^{-1}(T) = k(T)$ .*

Of course, this bijection generalizes to a bijection between increasing  $G$ -connected forests with  $q$  components and broken circuit free subforests of  $G$  with  $q$  components.

## References

- [1] Norman Biggs. *Algebraic graph theory*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 1993.
- [2] David D. Gebhard and Bruce E. Sagan. A chromatic symmetric function in noncommuting variables. *J. Algebraic Combin.*, 13(3):227–255, 2001.
- [3] Ira M. Gessel. Enumerative applications of a decomposition for graphs and digraphs. *Discrete Math.*, 139(1-3):257–271, 1995. Formal power series and algebraic combinatorics (Montreal, PQ, 1992).
- [4] Ira M. Gessel and Bruce E. Sagan. The Tutte polynomial of a graph, depth-first search, and simplicial complex partitions. *Electron. J. Combin.*, 3(2):Research Paper 9, approx. 36 pp. (electronic), 1996. The Foata Festschrift.
- [5] Richard P. Stanley. A symmetric function generalization of the chromatic polynomial of a graph. *Adv. Math.*, 111(1):166–194, 1995.

- [6] Richard P. Stanley. Graph colorings and related symmetric functions: ideas and applications: a description of results, interesting applications, & notable open problems. *Discrete Math.*, 193(1-3):267–286, 1998. Selected papers in honor of Adriano Garsia (Taormina, 1994).
- [7] Hassler Whitney. A logical expansion in mathematics. *Bull. Amer. Math. Soc.*, 38:572–579, 1932.